



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$4s = 8\pi(1 + \frac{1}{3} - \frac{1}{18} - \frac{1}{729} - \frac{1}{1048576} - \frac{875}{3888} - \frac{7}{478296} - \frac{847}{1190042784} - \dots).$$

$$4s = 26.0104 \text{ exact to two decimal places.}$$

Also solved by F. P. Matz, and J. F. W. Scheffer.

PROBLEMS.

30. Proposed by R. J. ADCOCK, Larchland, Illinois.

When the sum, of the distances of a point of a plane surface, from all its other points, is a minimum, that point is the centre of gravity of the plane surface.

QUERIES AND INFORMATION.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

"The Mysterious Formula."

Referring to my article on "Logarithms of Negative Numbers" published in the April Number, Vol. I., Mr. C. D. Schmitt, on page 214, deduces the following singular result: $\pi = \frac{\log(-1)}{\sqrt{-1}}$.

Another very remarkable result can be deduced from this as follows:

Dividing by 2, we have $\frac{1}{2\sqrt{-1}} \log(-1) = \frac{1}{2}\pi$. This may be written $\log \sqrt{-1} = \frac{1}{2}\pi$. $\therefore \sqrt{-1} = e^{\frac{1}{2}\pi} = 4.810477381$. This is what Professor Benjamin Peirce in his linear Associative Algebra, p. 5 (edition published by D. Van Nostrand), calls "the mysterious formula."

Writing i for $\sqrt{-1}$, the formula is $\frac{1}{i} = \sqrt{e^\pi}$.

M. C. STEVENS.

Editor Finkel has notified me that he cannot spare the space for further discussion of the possibility of a root of the equation

$4 + \sqrt{x-4} - \sqrt{x+4} = 0$. I will therefore again refer Mr. Draughton to Mr. Horner's article in the Philosophical Magazine, and also to Wentworth's Algebraic Analysis, page 278 to page 286.

H. C. WHITAKER.

The International Mathematical Congress.

Professor A. Vasiliev, President of the Physico-mathematical Society